

UNCLASSIFIED

AD NUMBER
ADB259894
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Jul 99. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St., Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 17 Jun 2002

THIS PAGE IS UNCLASSIFIED

AD _____

GRANT NUMBER DAMD17-98-1-8075

TITLE: The Essential Role of Protein Phosphatase-1 in Mitogenic Signaling and Breast Cancer

PRINCIPAL INVESTIGATOR: Carey J. Oliver

CONTRACTING ORGANIZATION: Duke University Medical Center
Durham, North Carolina 27710

REPORT DATE: July 1999

TYPE OF REPORT: Annual Summary

PREPARED FOR:

U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, July 1999). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 4

20001121 094

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND


Award Number: DAMD17-98-1-8075


Organization: Duke University Medical Center

Location of Limited Rights Data (Pages):

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.





REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE July 1999 3. REPORT TYPE AND DATES COVERED (1 Jul 98 - 30 Jun 99) Annual Summary

4. TITLE AND SUBTITLE The Essential Role of Protein Phosphatase-1 in Mitogenic Signaling and Breast Cancer 5. FUNDING NUMBERS DAMD17-98-1-8075

6. AUTHOR(S)
Carey J. Oliver

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)
Duke University Medical Center
Durham, North Carolina 27710

8. PERFORMING ORGANIZATION
REPORT NUMBER

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)
U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

10. SPONSORING / MONITORING
AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT

Distribution authorized to U.S. Government agencies only (proprietary information, July 1999). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)
There is growing evidence that protein dephosphorylation is as important as protein phosphorylation in cell proliferation. I am currently studying the role of protein phosphatase 1 (PP1), a serine/threonine phosphatase, in mitogenic signaling and breast cancer progression. To accomplish this goal, the following objectives are being addressed: (1) Determination of the changes in expression and function of MAP kinase and PP1 isoforms in a diversity of breast cancer cells. (2) Using *Xenopus* oocytes, we are delineating the role of PP1 in the activation of the MAP kinase pathway. (3) Identification of regulatory proteins capable of binding to PP1 and modulating its function in mitogenic signaling by screening *Xenopus* and MCF-7 expression libraries. (4) Determination of the effect of PP1 regulatory proteins on its function in mitogenic signaling and growth regulation. We have begun work on objectives one and two and made significant progress on objective three. We have identified a number of *Xenopus* proteins which bind to PP1 and are focusing on two proteins, spinophilin and PKC Zeta. We have begun *in vivo* studies of the interactions between PP1 and these two proteins and shown that they can each be co-immunoprecipitated with PP1. Future work will further delineate the biological consequences of PP1's interaction with these two proteins.

14. SUBJECT TERMS
Breast Cancer, Protein Phosphatase-1, MAP Kinase, signal transduction, *Xenopus laevis*

15. NUMBER OF PAGES
10

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT
Unclassified

18. SECURITY CLASSIFICATION OF THIS
PAGE
Unclassified

19. SECURITY CLASSIFICATION
OF ABSTRACT
Unclassified

20. LIMITATION OF ABSTRACT
Limited

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

CJO Where copyrighted material is quoted, permission has been obtained to use such material.

CSB Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

CJO Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

NA In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

n/a For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

NA In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

NA In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

NA In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

PI - Signature _____ Date _____

Table of Contents

A. Front Cover -	page 1
B. Standard Form 298	page 2
C. Foreward	page 3
D. Table of Contents	page 4
E. Annual Summary	pages 5-8
F. Key Research Accomplishments	page 9

Annual Summary

The role of protein phosphorylation in mitogenic signaling is well established. Much of the focus of this research has been on protein kinases, while the importance of protein phosphatases has not been addressed to the same extent. There is growing evidence that protein dephosphorylation is as important as protein phosphorylation in cell proliferation. I am currently studying the role of protein phosphatase 1 (PP1), a serine/threonine phosphatase, in mitogenic signaling and breast cancer progression. To accomplish this goal, the following objectives are being addressed: (1) Determination of the changes in expression and function of MAP kinase and PP1 isoforms in a diversity of breast cancer cells. (2) Using *Xenopus* oocytes, a model system for the study of cell cycle regulation and the MAP kinase pathway, we are delineating the role of PP1 in the activation of the MAP kinase pathway. (3) Screening *Xenopus* and MCF-7 expression libraries to identify regulatory proteins capable of binding to PP1 and modulating its function in mitogenic signaling. (4) Determination of the effect of PP1 regulatory proteins on its function in mitogenic signaling and growth regulation, particularly in MCF-7 breast cancer cells. The completion of this study will define the role of PP1 and its regulators in mitogenic signaling and their potential contribution to breast cancer progression.

Objective one – Levels of the four different PP1 isoforms in MCF-7 cells were established by western blot using isoform specific antibodies obtained from Dr. Brian Wadzinski. MCF-7 cells predominately contain the α isoform of PP1, with lesser amounts of the β and $\gamma 1$ isoforms, and no appreciable amount of the $\gamma 2$ isoform (data not shown). Next, the total levels of PP1 were checked in several different breast cancer cell lines by western blot using a non-isoform specific PP1 antibody. It was determined that the total level of PP1 was similar in SKBR3, MD468, MCF-7, HBL100, TH7d, ZR75-1, and a control, noncancerous breast epithelial cell line, NC26 (data not shown). However, it is possible that one or more isoforms of PP1 has varied expression levels in breast cancer cells. Future work will determine whether the levels and activity of PP1 isoforms and MAPK change in the various breast cancer cell lines when compared to control cell lines.

Objective two – The initial observation that Inhibitor-1, a specific endogenous inhibitor of PP1, prevents Mos activation of the MAP kinase cascade in *Xenopus* oocyte extracts has been difficult to repeat because of the extreme variability in *Xenopus* oocyte extracts. Therefore, we have temporarily put aside these experiments and concentrated on objective three, which has produced very interesting results. Work on objective two will be continued in the near future.

Objective three – We have performed an expression screen of a *Xenopus* ovary phage cDNA library, kindly provided by Dr. Sally Kornbluth, using digoxigenin labeled PP1. Briefly, bacteria were infected with the phage library and grown on agar plates until plaques formation began. Nitrocellulose filters were soaked in IPTG and placed on the agar plates to induce protein expression in the phage-infected bacteria. The filters were then incubated with digoxigenin-labeled PP1 and subsequently with an I^{125} labeled anti-digoxigenin antibody. Then the filters were visualized using autoradiography and positive plaques were cored and amplified. After performing two sets of three rounds of

screening, we have identified 37 positive clones which encode 23 different proteins (See Table 1).

Clone # isolates Identity			Clone # isolates Identity		
XPBP1	2	putative glucoamylase (Gm?)	XPBP13	1	Novel
XPBP2	3	putative ATP binding protein	XPBP14	1	Novel
XPBP3	3	PKC Zeta	XPBP15	2	Ral A
XPBP4	2	Novel	XPBP16	1	Putative Translational Release Factor
XPBP5	7	Neurabin II/Spinophilin	XPBP17	1	Novel
XPBP6	1	Zona pellucida B	XPBP18	1	No Insert
XPBP7	1	Novel	XPBP19	1	Putative 60S ribosomal protein L18A
XPBP8	1	putative adipose differentiation related protein	XPBP20	2	Threonyl-tRNA Synthetase
XPBP9	1	Putative BTG family protein	XPBP21	1	Glutamine Synthetase
XPBP10	1	Putative ASMTL	XPBP22	1	Putative Tropomyosin
XPBP11	1	Putative Nucleosome Assembly Protein	XPBP23	1	Putative Caspase
XPBP12	1	RLF			

Table 1 – Clones isolated from expression screen of *Xenopus* ovary phage cDNA library.

Once the 5' ends of the clones were sequenced, we wanted to confirm that these proteins were in fact PP1 binding proteins. First, we attempted to bacterially express the clones directly from the library vector. Several attempts were unsuccessful at producing any protein. We then produced S³⁵ labeled protein using an *in vitro* transcription/translation reaction to use in *in vitro* binding assays. Unfortunately, the background in these assays was unacceptably high. Therefore, we decided to concentrate on two known proteins, clone 3 and clone 5.

Clone 3 encodes for amino acids 49-510 of the atypical protein kinase C isoform zeta which had been previously cloned in *Xenopus*. The atypical PKCs, unlike the classical PKC isoforms, are not activated by calcium or diacylglycerol, but by lipids. Interestingly, it has already been shown that PKC Zeta can activate the MAP kinase pathway in *Xenopus* oocytes and cause maturation in the absence of hormonal stimulus (1). Thus, it is exciting to hypothesize that the mechanism of action of PP1 in oocyte maturation is to activate PKC Zeta and subsequently the MAP kinase pathway. However, it is unknown whether or not PKC Zeta is in the same pathway as Mos. Our future research will address this question. To further delineate the PKC Zeta/PP1 interaction we have obtained several myc-tagged PKC Zeta constructs from Dr. Peter Parker and expressed them in HEK-293 cells to look at the *in vivo* interaction with PP1. There are two sites on PKC Zeta which are especially important for its activation, the pseudosubstrate domain and Threonine-410. Threonine-410 is in the activation loop of

the kinase domain and the protein is activated when phosphorylated on this residue. This phosphorylation releases the pseudosubstrate domain from the catalytic site allowing PKC Zeta to phosphorylate its substrates. We have shown that endogenous PP1 can co-immunoprecipitate wild-type and constitutively active mutants of PKC Zeta (A119E or Δ PSS), but not kinase dead PKC Zeta mutants(T410A) (Figure 1).




Figure 1 – Co-immunoprecipitation of PP1 and myc-PKC Zeta from HEK-293 cells. Briefly, various myc-tagged PKC Zeta constructs were transiently overexpressed in HEK-293 cells, the cells were lysed, and immunoprecipitated with either PP1 or myc antibodies. Samples were run on SDS-PAGE, transferred to nitrocellulose, and blotted with anti-myc antibody. The higher molecular weight band is PKC Zeta.

In addition, PKC Zeta can not be isolated from HEK-293 cell lysates using microcystin sepharose, a small molecular weight marine toxin which binds to the catalytic site of PP1 (data not shown). This suggests that PKC Zeta may be a substrate of PP1. In support of this idea, a recent paper has shown that okadaic acid, a PP1 and PP2A inhibitor, can activate PKC Zeta in 3T3 L1 adipocytes (2). Future research will further study the nature of this interaction and the biological consequences of it.

Clone 5 encodes for amino acids 1-719 of spinophilin/neurabin II, a previously identified PP1 binding protein which is ubiquitously expressed in mammalian tissues. Spinophilin contains an N-terminal actin-binding domain followed by a PP1 binding motif. It also has a PDZ domain and a C-terminal coiled-coil motif. This protein has not previously been cloned from *Xenopus*. Thus, we have learned from a sequence alignment of the rat and frog proteins that the PP1 and PDZ domains are highly homologous, indicating that these regions are very important for spinophilin function. In addition, we have been able to co-immunoprecipitate endogenous PP1 with endogenous spinophilin. We have also been able to purify spinophilin using microcystin sepharose indicating that it is not a substrate for PP1 (data not shown). We have also shown that spinophilin is a phosphoprotein *in vivo* and a PKA substrate *in vitro* (data not shown). How this phosphorylation affects PP1 binding and spinophilin function inside a cell will be a subject of future study. In addition, we will attempt to determine spinophilin's effect on oocyte maturation by microinjection of antisense oligonucleotides or peptides which disrupt PP1 binding.

Objective Four – work has not yet begun on objective four.

References

1. Berra,E., Diaz-Meco,M.T., Dominguez,I., Municio,M.M., Sanz,L., Lorzano,J., Chapkin,R.S. and Morcat,J. (1993). *Cell* 74, 555-63.
2. Standaert,M.L., Bandyopadhyay,G., Sajan,M.P., Cong,L., Qwon,M.J. and Farese,R.V. (1999). *J. Biol. Chem.* 274, 14074-78.

Key Research Accomplishments

- Determined levels of different PP1 isoforms in MCF-7 cells
- Determined total levels of PP1 in several different breast cancer cell lines
- Screened a *Xenopus* ovary phage cDNA library using digoxigenin-labeled PP1 as a probe. Focused attention on 2 clones for further research
- Showed association of PP1 and PKC Zeta, as well as, PP1 and spinophilin in HEK-293 cells by co-immunoprecipitation
- Showed that spinophilin could be isolated using microcystin sepharose and PKC Zeta could not be isolated using microcystin sepharose
- Established that spinophilin is a phosphoprotein *in vivo* and a PKA substrate *in vitro*



DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

17 Jun 02

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218

SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for grants. Request the limited distribution statements for the Accession Documents listed at enclosure be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Judy Pawlus at DSN 343-7322 or by e-mail at judy.pawlus@det.amedd.army.mil.

FOR THE COMMANDER:

Encl

PHYLLIS M. PINEHART
Deputy Chief of Staff for
Information Management

ACCESSION DOCUMENT NUMBERS

ADB266028

ADB259887

ADB259894

ADB259837

ADB258947

ADB274390

ADB262488

ADB257222

ADB274382

ADB258931

~~ADB260157~~

~~ADB274457~~

> Do not downgrade